GENERAL INDEX TO VOLUME XI

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italics*; and previously published scientific names and all other matter, in ordinary type.

A

acerinus (Aleurodiscus), 40
Acidity, relation of, to growth of certain wood-destroying fungi, 43
Actinostroma, 1; crassum, 2
Agar culture solutions, 52
albido-carneum (Asterostroma), 29
albus (Microstroma), 27
Aleurodiscus acerinus, 40
americanorum (Microstroma), 27
ampla (Auriculariopsis), 10
ampla (Cyphella), 9
anomala (Peziza), 19
anomala (Solenia), 19
anomala var. ochracea (Solenia), 19
anomaloides (Solenia), 19
applanatus (Fomes (Ganoderma)), 37
Armillaria mellea, growth of, in relation to H-ion concentration of media, 47
Asterostroma, 28; albido-carneum, 29; bicolor, 32; cervicolor, 25; corticola, 28; gracile, 34; muscicolum, 31; ochrostroma, 34; pallidum, 29; spiniferum, 33
Auriculariopsis, 9; ampla, 10

B

Beccariella, 1
bicolor (Asterostroma), 32
Burt, E. A. Some wood-destroying
fungi of Java, 37; The Thelephoraceae
of North America, XIII, 1

C

candida (Solenia), 14
Candolleana (Cladoderris), 2
Celluloses in nutrient solutions, 50
cervicolor (Asterostroma) 28
cervicolor (Corticium), 28
cinerea (Solenia), 22
Cladoderris, 1; Candolleana, 2; crassa, 2;
dendritica, 2, 36; floridana, 4, 36;
infundibuliformis, 3
conferta (Solenia), 17
confusa (Solenia), 19
corrugata (Trametes), 39

ANN. Mo. Bot. GARD., Vol. 11, 1924

Corticium cervicolor, 28; flocculentum, 9; salicinum, 10 corticola (Asterostroma), 28 crassa (Cladoderris), 2 crassum (Actinostroma), 2 Craterellus marasmioides, 7; pulverulentus, 7; spathularius, 8 Cymatella, 6; minima, 6, 36; marasmioides, 7; pulverulenta, 7, 36 Cymatolerma, 1 Cynorrhynchium, 124 Cyphella ampla, 9; endophila, 25 Cytidia, 9; flocculenta, 9; rutilans, 10, 36; salicina, 10; tremellosa, 12, 36

D

Daedalea (Peziza), 23
Daedalea confragosa, growth of, in relation to H-ion concentration of media, 47
dendritica (Cladoderris), 2, 36
Diplacus, 124, 326; arachnoideus, 328; aridus, 336; aurantius, 338; calycinus, 331; cardinalis, 138; Clevelandii, 327; glutinosus, 338, var. aurantiacus, 338, var. Godroni, 349, var. grandiflorus, 335, var. latifolius, 339, var. puniceus, 342, var. splendidus, 349, var. stellatus, 337, var. Verschaffeltii, 349; X Godroni, 349; grandiflorus, 335; latifolius, 338; leptanthus, 334, 335; linearis, 334; longiflorus 328, 335; parviflorus, 344; puniceus, 344; rugosus, 350; speciosus, 328; × splendidus, 349; stellatus, 337; × Verschaffeltii, 349
Dodge, C. W., Zeller, S.M. and. Leucogaster and Leucophlebs in North America, 389

E

elongatus (Polystictus), 37 Endogone Tozziana 403 endophila (Cyphella), 25 endophila (Solenia), 25 Erythranthe, 124, 137; cardinalis, 138 Eumimulus, 126, 137

(461)

Eunanus, 268
Eunanus, 268; angustatus, 315; angustifolius, 298; Austinae, 296; bicolor, 307; Bigelovii, 277; Bolanderi, 273; brevipes, 270; Breveri, 261; clivicola, 297; Congdonii, 311; Couteri, 313; Cusickii, 282; Douglasii, 319, 321, var. parviflorus, 311; Fremontii, 283, 289; Kelloogii, 318, var. parviflorus, 311; latifolius, 310; Layneae, 291; leptaleus, 304; mephiticus, 300; mohavensis, 308; nanus, 295, a pluriflorus, 294; Parryi, 276; pictus, 324; pulchellus, 316; Rattanii, 289; subsecundus, 285; subuniflorus, 321; Tolmei, 294; Torreyi, 289; tricolor, 313

fasciculata (Solenia), 15 filicina (Solenia), 18 Filter-paper solutions, 51 flocculenta (Cytidia), 9
flocculenta (Thelephora), 9 flocculenta (Thelephora), 9
flocculentum (Corvicium), 9
floridana (Cladoderris), 4, 36
foetidus (Hypolyssus), 5
Fomes (Ganoderma) applanatus, 37;
elongatus, 37; Korthalsii, 37; pectinatus, 37; velutinosus, 37
Fungi: growth of certain wood-destroying, in relation to the H-ion concentration of the media 431; some wood-

tration of the media, 431; some wood-destroying, of Java, 37; studies in the physiology of the, XVII, 43

G

gracile (Asterostroma), 34 gracilis (Solenia), 26 gracilis (Solenia), 15 Grant, Adele Lewis, A monograph of the genus Mimulus, 99

H

Harris, J. Arthur. Variation and correlation in the inflorescence of Manfreda virginica, 411 Heterochaete tenuicula, 40 Hydnaceae, 39 Hydnangium liospermum, 407; nudum, Hydnum obrutans, 39, 42

Hydrogen-ion concentration of the me-dia, growth of certain wood-destroy-ing fungi in relation to the, 43 Hymenochaete muscicola, 31; nigricans,

Hymenogaster anomalus, 399

Hypolyssus, 4; foetidus, 5; Montagnei, 5.

I

infundibuliformis (Cladoderris), 3 ingainicola (Microstroma), 27

Java, some wood-destroying fungi of, 37 javensis (Protomerulius), 41, 42
Jefferson Barracks, Mo., type and variation of Manfreda virginica at, 415 Juglandis (Microstroma), 27

Korthalsii (Fomes), 37

L

Lenzites sepiaria, growth of, in relation to the H-ion concentration of the media, 47

Leucogaster and Leucophlebs in North

America, 389
Leucogaster, 389; sp., 403; anomalus, 399, 410; araneosus, 399, 410; badius, 400, 410; Bucholtzii, 404; citrinus, 398, 410; floccosus, 402, 410; foveolatus, 397, 410; fragrans, 403; fulvimaculosus, 401, 410; liosporus, 402; luteo-maculatus, 394, 410; nudus, 404; odoratus, 396, 410; rubescens, 395, 410; Toxsians, 403, 410
Leucophlebs, 405; gibbosum, 397; candids, 407, 410; magnata, 406, 410
Leucophleps, 389; citrina, 398; foveolata, 397; odorata, 396 America, 389

397; odorata, 396 leucosporum (Microstroma), 27

Lomatia, 9; salicina, 10

M

marasmioides (Craterellus), 7 marasmioides (Cymatella), 7 Manfreda virginica, variation and cor-relation in the inflorescence of, 411 Matruchotia, 26; varians, 26 Media: growth of certain wood-destroying fungi in relation to the H-ion concentration of the, 43, 48; peptone-nutrient solution with sugar, 48, without sugar, 48; Richards' E solution, 48

medulla-panis (Poria), 38 Meramec Highlands, Mo., type and variation of Manfreda virginica at, 415

Microstroma, 26; albus, 27; american-orum, 27; ingainicola, 27; Juglandis, 27; leucosporum, 27

Mimulastrum, 308

Mimulus, 124; acaulis, 151; acutangulus, 124; acaulis, 151; acutangulus, 127; acutidens, 202; acutidens, 186, 203; alatus, 131; alatus f. albiforus, 132; albus, 345; Alecterolophus, 350; alpinus, 154; alsinoides, 232, var. minimus, 233; alsinoides β paniculatus, 232; ampliatus, 214; andicolus, 188, 232; ampliatus, 214; andicolus, 232; ampliatus, 214; andicolus, 232; ampliatus, 214; andicolus, 232; ampliatus, 234; andicolus, 234; ampliatus, 234; ampliatus, 234; andicolus, 234; ampliatus, 232; ampliatus, 214; andicolus, 188; androsaceus, 253; angustatus, 315; angustatus, 316; angustifolius, 298; angustatus, 316; angustifolius, 298; angustifolius, 135; arenarius, 215, 384; aridus, 336; arvensis, 174; asamicus, 206; atropurpureus, 321; aurantiacus, 338; aurantiacus, 146; Austinae, 296; Bakeri, 177; burbatus, 260; K Bartonianus, 347; bicolor, 247, 388; Bigelovii, 277, var. cuspidatus, 279; Bigelovii var. ovatus, 252; Bioletti, 249; Bodinieri, 208; Bolanderi, 273, var. brachvdontus, 275. Bioletti, 249; Bodinieri, 208; Bolanderi, 273, var. brachydontus, 275; breviforus, 200; brevipes, 270; brevipes, 273; Breweri, 261, 386; Bridgesii, 210, var. integrifolia, 211, var. stolonifera, 211; caespitosus, 154; cardinalis, 138; cardinalis, 242, var. exsul, 138, var. griseus, 138, var. rigens, 138; Clarkii, 315; clementinus, 150; Clavalendii, 227; divisola, 207. 159; Clevelandii, 327; clivicola, 297; coccineus, 302; Colensoi, 197; Congdonii, 311; Congdonii, 317; cornalionus, 155; cordatus, 177; × cornation, 347; Coulteri var. angustatus, 315; crinitus, 186; Crista-galli, 350; cupreus, 150; Cusickii, 281; cuspidatus, 177; debitis, 137; decurtatus, 288; deflexus, 260; deltoideus, 216; densus, 298; dentatus, 209; dentatus var. gracilis, dentatus, 209; dentatus var. gracilis, 226; dentilobus, 185; depressus, 186, var. nanus, 187; depressus var. acaulis, 151, var. Pissisi, 346; diffusus 254, 578; discolor, 257; Douglasii, 320; Douglasii, 318; Dudleyi, 235, 376; X duplex, 347; Eastwoodiae, 242; Eisenii, 183; X Elphinstonea, 347; equinus, 159; erosus, 177; exiguus, 256; exilis, 350; filicaulis, 251; floribundus, 216, var. geniculatus, 220, var. membranaceus, 221, var. subulatus, 222; floribundus, 221, var. subulatus, 222; floribundus tus, 220, var. membranaceus, 221, var. subulatus, 222; floribundus β minor, 216; formosana, 206; × Forsythiana, 347; Fremontii, 283; geniculatus, 220; Geyeri, 190; glabratus, 188, var. Fremontii, 190, var. parvi-Horus, 194; glabratus var. ascendens, 158, var. Jamesti, 191; glandulosus, 345; glareosus, 177; glaucescens, 175; glutinosus, 338, var. brachypus, 328, var. linearis, 334, var. puniceus, 342; gracilipes, 252; gracilis, 134; grandis, 159; grandiflorus, 159; gratioloides, 266; Grayi, 203, 380; guttatus, 157, var. arvensis, 174, var. decorus, 173, var. depauperatus, 170, var. Hallii, 172, var. puberulus, 170; guttatus, 148, var. grandis, 158, var. insignis, 181; Hallii, 172; × Harrisonia, 347; hirsutus, 159, 350; × hybridus, 348; implexus, 151, 155; implicatus, 151; inamoenus, 191; inconspicuus var. acutidens, 202, var. latidens, 201; inodorus, 202, var. latidens, 201; inodorus, 203; Jamesti, 190, var. Fremontirus, 230; Jamesii, 190, var. Fremontii, 190, var. texensis, 188; javanicus, 350; Jepsonii, 306; Johnstonii, 280; jun-germannioides, 222; Kelloggii, 318, Jepsonii, 306; Johnstonii, 280; jungermannioides, 222; Kelloggii, 318, 388; Kingii, 194; laciniatus, 183; lanatus, 345; Langsdorfii, 158, 177, var. alpinus, 151, var. argutus, 158, var. arvensis, 174, var. californicum, 159, var. grandis, 159, var. gutatus, 159, var. insignis, 181, var. microphyllus, 171, var. minor, 151, var. nasutus, 177, var. platyphyllus, 158, var. Tilingi, 155; latifolius, 310; latidens, 201; Layneae, 291, 386; Leibergii, 231, 380; leptaleus, 304; leptanthus, 335; Lewisii, 236, 384; Levisii var. alba, 237, var. exsertus, 237, var. tetonensis, 237, f. tetonensis, 237; linearis, 136, var. lutea, 137; linearis, 334; longiforus, 328, 588, var. calycinus, 331, var. linearis, 334, × puniceus, 332, var. rutilus, 333; longiese, 211; longulus, 171; var. calycinus, 331, var. linearis, 334; × puniceus, 332, var. rutilus, 333; longipes, 211; longulus, 171; lucens, 155; lugens, 142; luteus, 146, var. alpinus, 149, × var. calycanthemus, 348, var. nummularis, 149, × var. pardinus, 348, var. rivularis, 148, var. variegatus, 148, × var. Wilsoni, 348, var. Youngana, 148; luteus, 158, 177, var. alpinus, 151, 154, var. avarantivar. alpinus, 151, 154, var. auranti-acus, 146, var. cuprea, 150, var. de-pauperatus, 170, 177, var. gracilis, 158, subvar. macrophyllus, 146, var. micranthus, 194; lyratus, 158; × Maclainianus, 348; × maculatus, 348; × maculatus, 348; × maculosus, 348; madagascariensis, 136; madrensis, 191; marmoratus, 174; membranaceus, 221; mephiticus, 300; membranaceus, 221; mephiticus, 300; micranthus, 182; microphyllus, 170; minor, 151; minhodes, 131; minusculus, 177; modestus, 317; mohavensis, 308; monitiformis, 226; monticides, 258; monticides, 264; moschatus, 223, var. longiflorus, 226, 386, var. sessilifolius, 229; moschatus, 226, var. pallidiflorus, 226; nanus, 294; nanus, 187, var. bicolor, 307, α pluriflorus, 294, β subuniflorus, 321; nasutus, 177, 383, var. insignis, 181, var. micranthus, 182; Nelsonii, 144; nemicranthus, 182; Nelsonii, 144; nepalensis, 206, var. japonica, 207, var. procerus, 207; Neubertii, 346; nevadensis, 243; nudatus, 176; numularis, 149; ocellatus, 148; orbicularis, 198; orizabae, 346; pachystylus, 234; pallens, 185; pallidus, 127; Palmeri, 250; Palmeri var. androsaceus, 253; paniculatus, 159; Parishii, 235; Parishii, 177; Parryi, 276; parviflorus, 344; parviflorus, 194, \$\beta\$ Bridgesis, 210; parvulus, 185; peduncularis, 213, 216; perfoliatus, 350; pietus, 324; Pilingi, 151; pilosellus, 243; pilosiusculus, 187; pilosus, 350; Pissisi, 346; Prattenii, 247; prionophyllus, 159; primuloides var. pilosellus, 243; propinquus, 194; prostratus, 199; pteropus, 346; puberulus, 170, 177; pubescens, 216; pulchellus, 316; Pulsiferae, 211, 388; punctatus, 346; purberulus, 195; pursillus, 135; purpureus, 255, 378, var. paurillus, 255; pusillus, 199; pusillus, 135; pygmaeus, 312; quinquevulnerus, 346; radicans, 350; Rattani, 288; reniformis, 191; repens, 197; ringens, 127, var. minthodes, 131; ringens var. congesta, 127, f. Peckii, tani, 288; reniformis, 191; repens, 197; ringens, 127, var. minthodes, 131; ringens var. congesta, 127, f. Peckii, 128; rivularis, 158; Roezli, 151; roseus, 237, var. glabrior, 237, var. Maclainianus, 348; rubellus, 266; rubellus, 259, 261, var. latiflorus, 259; × rubiginosus, 348; rupestris, 142; × sanguineus, 348; saxorum, 350; Scouleri, 158, var. caespitosus, 154; serotinus, 216; sessilifolius, 208; × Seymouriana, 349; × Smithii, 349; Smithii, 149; spissus, 277, 382; stamineus, 302; stellatus, 337; striatus, 135; strictus, spistus, 211, 302; stainingus, 302, stellatus, 337; striatus, 135; strictus, 134; subreniformis, 177; subsecundus, 285, 388, var. constrictus, 287, var. viscidus, 286; subuniforus, 321; Suksviscidus, 286; subuniflorus, 321; Suksdorfii, 264; subuniflorus, 187; tenellus, 171, 206; tener, 188; thermalis, 171; X tigridioides, 349; Tilingi, 151, var. caespitosus, 154, var. corallinus, 155; Torreyi, 289; Traskiae, 323; Treleasel, 325, 374; tricolor, 313; tricolor var. angustatus, 315; Uvedaliae, 136; variegatus, 148; verbenaceus, 142; veronicifolius, 155; violaceus, 350; viscidus, 286; viscosus, 338; washingtonensis, 213; Whipplei, 184; Whitneyi, 307; × Youngii, 349 minima (Cymatella), 6, 36

Monavia, 124 Montagnei (Hypolyssus), 5, 36 muscicola (Hymenochaete), 31 muscicolum (Asterostroma), 31 N

nigricans (Hymenochaete), 40 nigricans (Protocoronospora), 27 nothopus (Polystictus), 38, 42 notopus (Polyporus), 38

obrutans (Hydnum), 39, 42 obscurans (Stereum), 39, 42 ochracea (Solenia), 19 ochrostroma (Asterostroma), 34 Octaviania liosperma, 407; (Gautieria?) pityophila, 402; silesiaca, 402 Oenoe, 309

pallidum (Asterostroma), 29

Paradanthus, 195
pectinatus (Fomes), 37
Peptone-nutrient solution, 49
(Polystictus), 39 Persoonii (Polystictus), 39 Peziza anomala, 19; Daedalea, 23; poriae formis, 23; pruinata, 23; stipata, 19; tephrosia, 23 Pholiota adiposa, growth of, in relation to the H-ion concentration of the Pleurotus ostreatus, growth of, in relation to H-ion concentration of the media, 47 Polyporaceae, 37 polyporoidea (Solenia), 16 Polyporus adustus, growth of, in relation to H-ion concentration of the media, Polyporus notopus, 38 Polystictus elongatus, 37; nothopus, 38, 42; Persoonii, 39; spadiceus, 38; versicolor, growth of, in relation to H-ion concentration of the media, 47 H-ion concentration of the media Poria medulla-panis, 38, sp., 39 poriaeformis (Pesiza), 23 poriaeformis (Solenia), 23 Protocoronospora, 27; nigricans, 27 Protomerulius javensis, 41, 42 presingle (Pesign), 23 Protomerunus javenusi, 41, 42 pruinata (Peziza), 23 Pseudoenoe, 323 pubescens (Stereum), 10 pulverulenta (Cymatella), 7, 36 pulverulentus (Craterellus), 7

R

Richards' E solution, 48 rutilans (Cytidia), 10

S

salicina (Cytidia), 10
salicina (Lomatia), 10
salicina (Thelephora), 10
salicinum (Corticium), 10
Schizoplacus, 268
Schizophyllum commune, growth of, in relation to the H-ion concentration of the media, 47
Simiolus, 145, 195
Skepperia, 8; spathularia, 8, 36
Solenia, 13; anomala, 19; anomala var.
ochracea, 19; anomaloides, 19; candida, 14; cinerea, 22; conferta, 17; confusa, 19; endophila, 25; fasciculata, 15; filicina, 18; gracilis, 26; gracilis, 15; ochracea, 19; polyporoidea, 16; poriaeformis, 23; poriaeformis, 23; subportaeformis, 24; sulphurea, 18; villosa, 26; villosa, 18, var. polyporoidea, 16
spadiceus (Polystictus), 38

spadiceus (Polystictus), 38 spathularia (Skepperia), 8, 36 spathularius (Craterellus), 8 spiniferum (Asterostroma), 33 Stereum obscurans, 39, 42; pubescens, 10

stipata (Peziza), 19 subporiaciormis (Solenia), 24 sulphurea (Solenia), 18 Synplacus, 126

T

Temperature, relation of, to growth of wood-destroying fungi, 90 tenuicula (Heterochaete), 40 tephrosia (Peziza), 23
Thelephora flocculenta, 9; salicina, 10
Thelephoraceae of North America, XIII,
1
Trametes corrugata, 39
tremellosa (Cytidia), 12, 36
Tropanthus, 324

U

Uvedalia, 124; linearis, 136

V

varians (Matruchotia), 26 Variation and correlation in the inflorescence of Manfreda virginica, 411 velutinosus (Fomes), 37 villosa (Solenia), 26 villosa (Solenia), 18, var. polpyporoidea, 16

W

Wolpert, F. S. Studies in the physiology of the fungi, XVII. The growth of certain wood-destroying fungi in relation to the H-ion concentration of the media, 43

Z

Zeller, S. M., and Dodge, C. W. Leucogaster and Leucophlebs in North America, 389